



3rd International Olympiad on Astronomy and Astrophysics

Theoretical Competition

Please read these instructions carefully:

1. Each student will receive problem sheets in English and/or in his/her native language.
2. The available time for answering theoretical problems is 5 hours. You will have 15 short problems (Theoretical Part 1, Problem 1 to 15), and 2 long problems (Theoretical Part 2, Problem 16 and 17).
3. Use only the pen that has been provided on your desk.
4. Do **Not** use the back side of your writing sheets. Write only inside the boxed area.
5. Yellow scratch papers are not considered in marking.
6. Begin answering each problem in separate sheet.
7. Fill in the boxes at the top of each sheet of your paper with your "country name", your "student code", "problem number", and total number of pages which is used to answer to that problem.
- 8. Write the final answer for each problem in the box, labeled "Answer Sheet".**
9. Starting and the end of the exam will be announced by ringing a bell.
10. The final answer in each question part must be accompanied by units, which should be in SI or appropriate units. 20% of the marks available for that part will be deducted for a correct answer without units.
11. The required numerical accuracy for the final answer depends on the number of significant figures given in the data values in the problem. 20% of the marks available for the final answer in each question part will be deducted for answers without required accuracy as given in the problem. Use the constant values exactly as given in the table of constants.
12. At the end of the exam put all papers, including scratch papers, inside the envelope and leave everything on your desk.

Table of Constants

(All constants are in SI)

Parameter	Symbol	Value
<i>Gravitational constant</i>	<i>G</i>	
<i>Plank constant</i>	<i>h</i>	$6.63 \times 10^{-34} \text{ J s}$
<i>Speed of light</i>	<i>c</i>	
<i>Solar Mass</i>		
<i>Solar radius</i>		
<i>Solar luminosity</i>		
<i>Apparent solar magnitude (V)</i>		-26.8
<i>Solar constant</i>		
<i>Mass of the Earth</i>		
<i>Radius of the Earth</i>		
<i>Mean density of the Earth</i>		$5 \times 10^3 \text{ kg m}^{-3}$
<i>Gravitational acceleration at sea level</i>	<i>g</i>	
<i>Tropical year</i>		365.24 days
<i>Sidereal year</i>		365.26 days
<i>Sidereal day</i>		86164 s
<i>Inclination of the equator with respect to the ecliptic</i>	<i>ϵ</i>	
<i>Parsec</i>	<i>pc</i>	
<i>Light year</i>		
<i>Astronomical Unit</i>	<i>AU</i>	
<i>Solar distance from the center of the Galaxy</i>		
<i>Hubble constant</i>	<i>H</i>	
<i>Mass of electron</i>		
<i>Mass of proton</i>		
<i>Central wavelength of V-band</i>	<i>λ</i>	
<i>Refraction of star light at horizon</i>		
	<i>π</i>	3.1416

Useful mathematical formula:



3rd International Olympiad on Astronomy and Astrophysics

Theoretical Competition

Short Problems

Short Problems: (10 points each)

Problem 1: Calculate the mean mass density for a super massive black hole with total mass of $1 \times 10^8 M_{\odot}$ inside the Schwarzschild radius.

Problem 2: Estimate the number of photons per second that arrive on our eye at $\lambda = 550 \text{ nm}$ (V-band) from a G2 main sequence star with apparent magnitude of $m = 6$ (the threshold of naked eye visibility). Assume the eye pupil diameter is 6 mm and all the radiation from this star is in $\lambda = 550 \text{ nm}$.

Problem 3: Estimate the radius of a planet that a man can escape its gravitation by jumping vertically. Assume density of the planet and the Earth are the same.

Problem 4: In a typical Persian architecture, on top of south side windows there is a structure called "Tabeshband" (shader), which controls sunlight in summer and winter. In summer when the Sun is high, Tabeshband prevents sunlight to enter rooms and keeps inside cooler. In the modern architecture it is verified that the Tabeshband saves about 20% of energy cost. Figure (1) shows a vertical section of this design at latitude of $36^{\circ}.0 \text{ N}$ with window and Tabeshband.

Using the parameters given in the figure, calculate the maximum width of the Tabeshband, " x ", and maximum height of the window, " h " in such a way that:

- No direct sunlight can enter to the room in the summer solstice at noon.
- The direct sunlight reaches the end of the room (indicated by the point **A** in the figure) in the winter solstice at noon.




Figure (1)

Problem 5: The Damavand Mountain is located at the North part of Iran, in south coast of Caspian Sea. Consider an observer standing on the Damavand mountaintop (latitude = $35^{\circ} 57' N$; longitude = $52^{\circ} 6' E$; altitude $5.6 \times 10^3 m$ from the mean sea level) and looking at the sky over the Caspian Sea. What is the minimum declination for a star, to be seen marginally circumpolar for this observer. Geodetic radius of the Earth at this latitude is $6370.8 km$. Surface level of the Caspian Sea is approximately equal to the mean sea level.

Problem 6: Derive a relation for the escape velocity of an object, launched from the center of a proto-star cloud. The cloud has uniform density with the mass of M and radius R . Ignore collisions between the particles of the cloud and the launched object. If the object were allowed to fall freely from the surface, it would reach the center with a velocity equal to $\sqrt{\frac{GM}{R}}$.

Problem 7: A student tries to measure field of view (FOV) of the eyepiece of his/her telescope, using rotation of the Earth. To do this job, the observer points the telescope towards Vega (alpha Lyr., RA: 18.5^h , Dec: $+39^{\circ}$), turns off its "clock drive" and measures trace out time, $t=5.3$ minutes, that Vega crosses the full diameter of the FOV. What is the FOV of this telescope in arc-minutes?

Problem 8: Estimate the mass of a globular cluster with the radius of $r = 20 pc$ and root mean square velocity of stars equal to $v_{rms} = 3 kms^{-1}$.

Problem 9: The Galactic longitude of a star is $l = 15^{\circ}$. Its radial velocity with respect to the Sun is $V_r = 100 kms^{-1}$. Assume stars in the disk of the Galaxy are orbiting the center with a constant velocity of $V_0 = 250 kms^{-1}$ in circular orbits in the same sense in the galactic plane. Calculate distance of the star from the center of the Galaxy.

Problem 10: A main sequence star with the radius and mass of $R = 4R_{\odot}$, $M = 6M_{\odot}$ has an average magnetic field of $1 \times 10^{-4} T$. Calculate the strength of the magnetic field of the star when it evolves to a neutron star with the radius of $20 km$.

Problem 11: Assume the mass of neutrinos is $m_{\nu} = 10^{-5} m_e$. Calculate the number density of neutrinos (n_{ν}) needed to compensate the dark matter of the universe. Assume the universe is flat and 25 % of its mass is dark matter.

Hint: Take the classical total energy equal to zero

Problem 12: Calculate how much the radius of the Earth's orbit increases as a result of the Sun losing mass due to the thermo-nuclear reactions in its center in 100 years. Assume the Earth's orbit remains circular during this period.

Problem 13: Assume that you are living in the time of Copernicus and do not know anything about Kepler's laws. You might calculate Mars-Sun distance in the same way as he did. After accepting the revolutionary belief that all the planets are orbiting around the Sun, not around the Earth, you measure that the orbital period of Mars is 687 days, then you observe that 106 days after opposition of Mars, the planet appears in quadrature. Calculate Mars-Sun distance in astronomical unit (AU).

Problem 14: A satellite is orbiting around the Earth in a circular orbit in the plane of the equator. An observer in Tehran at the latitude of $\varphi = 35.6^\circ$ observes that the satellite has a zenith angle of $z = 46.0^\circ$, when it transits the local meridian. Calculate the distance of the satellite from the center of the Earth (in the Earth radius unit).

Problem 15: An eclipsing close binary system consists of two giant stars with the same sizes. As a result of mutual gravitational force, stars are deformed from perfect sphere to the prolate spheroid with $a = 2b$, where a and b are semi-major and semi-minor axes (the major axes are always co-linear). The inclination of the orbital plane to the plane of sky is 90° . Calculate the amplitude of light variation in magnitude (Δm) as a result of the orbital motion of two stars. Ignore temperature variation due to tidal deformation and limb darkening on the surface of the stars.

Hint: A prolate spheroid is a geometrical shape made by rotating of an ellipse around its major axis, like rugby ball or melon.